

<https://doi.org/10.37208/tgn27302>

Kames Bay, Isle of Cumbrae, Scotland: unexplored aspects of a Site of Special Scientific Interest

P.G. Moore

32 Marine Parade, Millport, Isle of Cumbrae
KA28 0EF

E-mail: p.geoff.moore@gmail.com

Kames Bay, Millport, Isle of Cumbrae, Scotland (Fig. 1) is one of the classic sandy beaches of the world. Probably more has been published on the identity, biology and distribution of macrofauna and meiofauna living there than on that of almost any other beach (King & Elmhirst, 1914; Stephen, 1928, 1929, 1931, 1932, 1938, 1945, 1953; Elmhirst, 1931, 1932; Nicholls, 1935, 1939; Watkin, 1938, 1939, 1941, 1942; Dixon, 1944; Clark & Milne, 1955; Smith, 1955; Ansell, 1961; Ansell & Trevallion, 1967; Mauchline, 1967; Ansell, 1969; Corey, 1969; McIntyre, 1970; Barnett, 1971; Ansell, 1972; Barnett & Watson, 1986; Powell & Moore, 1991; Moore & Beare, 1993; Beare & Moore, 1996, 1997, 1998; Cowie & Hannah, 2002, 2006; Moore, 2017).



Fig. 1. Kames Bay, Millport, Isle of Cumbrae, Scotland, 22nd March 2020, showing the glistening area of wetted sand on a receding tide revealed after two dry days but extensive rainfall in the previous weeks, with freshwater draining-up through the beach. (Photo: J. Moore)

For generations fieldwork classes have used this beach for practical and project work based at the Millport Marine Station (first under the Scottish Marine Biological Association (SMBA), then as the University Marine Biological Station Millport (UMBSM), and now as the Field Studies Centre Millport (FSCM)). Its designation as a Site of Special Scientific Interest (SSSI) was based on these educational considerations, but its uniqueness vis-à-vis its salinity conditions should guarantee such status. Intelligent decision-making about how best to manage SSSIs, and coasts in general, in a sensitive and sustainable way in order to conserve

biodiversity, requires basic science at the heart of an Integrated Coastal Zone Management (ICZM) policy.

The physical environment of Kames Bay has been known since the mid-1950s to be particularly interesting in terms of the freshwater upwelling that percolates through its upper regions (though this may occur to varying degrees down its whole extent). Rainfall from the island's catchment is channelled southwards along the underlying Great Cumbrae Fault – a geological feature that is responsible for the position of the embayment in the first place – until it surfaces through the beach (Smith, 1955). A thorough analysis of the physical environment of the beach, or its seasonal variability, has not been published (apart from extensive *in situ* temperature recordings by P.R.O. Barnett; see Barnett & Watson, 1986), although student classes have habitually sampled its physical properties.

Salinity is well known as a key environmental determinant in many aquatic ecosystems (Kinne, 1971), from rockpools (Daniel & Boyden, 1975; Morris & Taylor, 1983; McAllen *et al.*, 1998) to estuaries and lagoons (Barnes, 1996). However, little attention has been paid to the role of brackish water inside the beach at Kames Bay, which is needed because of its possible role in affecting macrofauna and other organisms on and within it. A clearer understanding of the salinity distribution in the beach aquifer would provide context for interpreting the long history of associated biological measurements.

Freshwater is often assumed to run-off beach surfaces with little impact on underlying salinity conditions in the sand (Reid, 1930; Hayward, 1994). Thus, Brown & McLachlan (1990) stated that “dramatic changes in beach water salinity seldom occur” (see also Barrett, 1974); but on Kames Bay, where brackish-water conditions prevail underground below Mean High Water Neap Tide (MHWN) level in an otherwise fully marine locality, they do (Smith, 1955; Hayward, 1994). Kames Bay thus merits particular consideration. As stressed by Eltringham (1971) “the work at Kames Bay ... shows that under certain unusual conditions, salinity can exert a limiting effect on the shore”. Large-scale perturbations in surface water salinity have been recorded in the water column off Millport (Thomason *et al.*, 1997), and such stochastic events with a high rate of change of salinity, may have a profound effect on the physiology, behaviour, distribution and abundance of marine organisms (Smith, 1955; Davenport *et al.*, 1975; Hayward, 1994; Barnes, 1996; Little & Kitching, 1996; Santos *et al.*, 1996).

Intermittent low salinity conditions may have a profound, and locally unrecognised, significance in terms of the behaviour and performance of beach-dwelling infauna, even to the extent of modifying productivity (Cowie & Hannah, 2002). Unpublished observations by P.R.O. Barnett (pers. comm.) made

offshore in Kames Bay after a period of sustained heavy rainfall some years ago revealed the sublittoral population of the infaunal bivalve *Fabulina* (previously *Tellina*) *fabula* all lying on top of the submerged sand. The suspicion is that groundwater pressure created inimical salinity/thixotropic conditions (in this case even extending to the immediate offshore), prompting their evacuation from the sand. Such a mechanism could result in whole populations of sensitive species being made vulnerable to transportation and/or predation, as also happens after some red tides (Griffith *et al.*, 2019). Groundwater discharge from the low-tide line has been shown to lead to the formation of algal blooms (Paerl, 1997). Many sandy beach-dwelling organisms, such as cockles (*Cerastoderma* spp.) and lugworms (*Arenicola* spp.), are economically important, so such data would not only be of academic interest.

The extent to which Kames Bay is unusual in terms of freshwater ingress interstitially is, as yet, to be established and it could well be that all pocket sandy beaches in hard landscape coves and embayments (and even extensive earth-backed sand strands) are generally subject to similar subterranean freshwater percolation influences after heavy rain.

Subterranean interfaces in littoral sediments also have a profound evolutionary significance. Brown & McLachlan (1990) noted that the discharge of groundwater through beaches, as unconfined aquifers, provides a path for the interstitial fauna to evolve landwards to terrestrial and freshwater habitats (Warwick, 1989). These authors note as well that towards the backshore of most sandy beaches a rapid change in interstitial salinity occurs, as percolating seawater gives way to inflowing groundwater. As Jansson (1967) has indicated, however, the importance of rain on interstitial salinity is mainly via increasing groundwater pressure (rather than by overflowing beaches). The up/downshore positioning of the interface between saline and fresh water influences within sandy beach aquifers and its temporal variability under different tidal regimes have been reported elsewhere (Robinson *et al.*, 2007; Heiss & Michael, 2014; McAllister *et al.*, 2015, Fig. 1) and these are known to drive microbial geochemical processes (Paerl, 1997; Cowie & Hannah, 2002; McAllister *et al.*, 2015). On Kames Bay, this universal interface is certainly slipped seawards, providing unique opportunities for research generating comparisons with estuarine interfaces (Phillips & Fleeger, 1985; Santos *et al.*, 1996; Robinson *et al.*, 2007). Cowie & Hannah (2002) associated the extremely low numbers of naked (non-testate) amoebae in the beach sand at Kames Bay with the unusual salinity regime there, potentially influencing nutrient recycling (McLachlan & Illenberger, 1986; Montagna & Yoon, 1991; Millham & Howes, 1994).

There have been few studies to date of salinity fluctuations within sandy beaches during tidal inundation, with one example being Heiss & Michael

(2014). Most published data that exist for salinity variations in sandy beaches (e.g. Jansson, 1966, 1967) have been derived at low tide. Of more than merely academic interest, such a study on Kames Bay would also assist in the formulation of soundly based advice to policy makers responsible for management and conservation decision-making, especially on sandy beach SSSIs. I hope that some enterprising marine scientist might be intrigued enough to research this topic locally.

ACKNOWLEDGEMENT

I am grateful to Professor James Heiss (Department of Environmental, Earth, and Atmospheric Sciences, University of Massachusetts Lowell, U.S.A.) for his most helpful comments that greatly improved the draft manuscript.

REFERENCES

- Ansell, A.D. (1961). Reproduction, growth and mortality of *Venus striatula* (Da Costa) in Kames Bay, Millport. *Journal of the Marine Biological Association of the United Kingdom* 41, 191–215. <https://doi.org/10.1017/S0025315400001648>
- Ansell, A.D. (1969). Association of the amphipod *Acidostoma neglectum* Dahl with the anthozoan *Peachia hastata* Gosse. *Journal of Natural History* 3, 345–347. <https://doi.org/10.1080/00222936900770301>
- Ansell, A.D. (1972). Distribution, growth and seasonal changes in biochemical composition for the bivalve *Donax vittatus* (da Costa) from Kames Bay, Millport. *Journal of Experimental Marine Biology and Ecology* 10, 137–150. [https://doi.org/10.1016/0022-0981\(72\)90099-8](https://doi.org/10.1016/0022-0981(72)90099-8)
- Ansell, A.D. & Trevallion, A. (1967). Studies on *Tellina tenuis* Da Costa. I. Seasonal growth and biochemical cycle. *Journal of Experimental Marine Biology and Ecology* 1, 220–235. [https://doi.org/10.1016/0022-0981\(67\)90016-0](https://doi.org/10.1016/0022-0981(67)90016-0)
- Barnes, R.S.K. (1996). *The Brackish-water Fauna of Northwestern Europe*. Cambridge University Press, Cambridge.
- Barnett, P.R.O. (1971). Some changes in intertidal sand communities due to thermal pollution. *Proceedings of the Royal Society of London B*, 177, 353–364. <https://doi.org/10.1098/rspb.1971.0035>
- Barnett, P.R.O. & Watson, J. (1986). Long-term changes in some benthic species in the Firth of Clyde, with particular reference to *Tellina tenuis* da Costa. *Proceedings of the Royal Society of Edinburgh* 90B, 287–302. <https://doi.org/10.1017/S0269727000005042>
- Barrett, J. (1974). *Life on the Sea Shore*. Collins, London.
- Beare, D.J. & Moore, P.G. (1996). The distribution, growth and reproduction of *Pontocrates arenarius* and *P. altamarinus* (Crustacea: Amphipoda) at Millport, Scotland. *Journal of the Marine Biological Association of the United Kingdom* 76, 931–950. <https://doi.org/10.1017/S0025315400040893>

- Beare, D.J. & Moore, P.G. (1997). The contribution of Amphipoda to the diet of certain inshore fish species in Kames Bay, Millport. *Journal of the Marine Biological Association of the United Kingdom* 77, 907–910.
<https://doi.org/10.1017/S0025315400036274>
- Beare, D.J. & Moore, P.G. (1998). Aspects of the life histories of *Perioculodes longimanus*, *Pontocrates arcticus* and *Synchelidium maculatum* (Crustacea: Amphipoda) at Millport, Scotland. *Journal of the Marine Biological Association of the United Kingdom* 78, 193–209.
<https://doi.org/10.1017/S0025315400040029>
- Brown, A.C. & McLachlan, A. (1990). *Ecology of Sandy Shores*. Elsevier Science Publishers, Amsterdam.
- Clark, R.B. & Milne, A.A. (1955). The sublittoral fauna of two sandy bays on the Isle of Cumbræ, Firth of Clyde. *Journal of the Marine Biological Association of the United Kingdom* 34, 161–180.
<https://doi.org/10.1017/S0025315400008663>
- Corey, S. (1969). The comparative life histories of three Cumacea (Crustacea): *Cumopsis goodsiri* (van Beneden), *Iphinoe trispinosa* (Goodsir) and *Pseudocuma longicornis* (Bate). *Canadian Journal of Zoology* 47, 695–704.
<https://doi.org/10.1139/z69-116>
- Cowie, P.R. & Hannah, F. (2002). Kames Bay – a sandy temperate shore inimical to naked amoebae. *32nd Annual Report of the University Marine Biological Station Millport*, 8-9.
- Cowie, P.R. & Hannah, F. (2006). Impact of hydrodynamic disturbance on the growth rates of marine naked amoebae. *Journal of the Marine Biological Association of the United Kingdom* 86, 39–42.
<https://doi.org/10.1017/S0025315406012835>
- Daniel, M.J. & Boyden, C.R. (1975). Diurnal variations in physico-chemical conditions within intertidal rockpools. *Field Studies* 4, 161–176.
- Davenport, J., Gruffydd, L.D. & Beaumont, A.R. (1975). An apparatus to supply water of fluctuating salinity and its use in a study of salinity tolerances of larvae of the scallop, *Pecten maximus* L. *Journal of the Marine Biological Association of the United Kingdom* 55, 391–409.
<https://doi.org/10.1017/S0025315400016015>
- Dixon, A.Y. (1944). Notes on certain aspects of the biology of *Cumopsis goodsiri* (van Beneden) and some other cumaceans in relation to their environment. *Journal of the Marine Biological Association of the United Kingdom* 26, 61–71.
<https://doi.org/10.1017/S0025315400014454>
- Elmhirst, R. (1931). Studies on the Scottish marine fauna. The Crustacea of the sandy and muddy areas of the tidal zone. *Proceedings of the Royal Society of Edinburgh* 51, 169–175.
<https://doi.org/10.1017/S0370164600023117>
- Elmhirst, R. (1932). Quantitative studies between tidemarks. *The Glasgow Naturalist* 10, 56–62.
- Eltringham, S.K. (1971). *Life in Mud and Sand*. English Universities Press, London.
- Griffith, A.W., Shumway, S.E. & Gobler, C.J. (2019). Differential mortality of North Atlantic bivalve molluscs during harmful algal blooms caused by the dinoflagellate, *Cochlodinium* (a.k.a. *Margalefidinium*) *polykrikoides*. *Estuaries and Coasts* 42, 190–203.
<https://doi.org/10.1007/s12237-018-0445-0>
- Hayward, P.J. (1994). *Animals of Sandy Shores*. Richmond Publishing Company, Slough.
- Heiss, J.W. & Michael, H.A. (2014). Saltwater-freshwater mixing dynamics in a sandy beach aquifer over tidal, spring-neap, and seasonal cycles. *Water Resources Research* 50, 6747–6766.
<https://doi.org/10.1002/2014WR015574>
- Jansson, B.-O. (1966). Microdistribution of factors and fauna in marine sandy beaches. *Veröffentlichungen des Instituts für Meeresforschung in Bremerhaven* 2, 77–86.
- Jansson, B.-O. (1967). Diurnal and annual variations of temperature and salinity of interstitial water in sandy beaches. *Ophelia* 4, 173–201.
<https://doi.org/10.1080/00785326.1967.10409619>
- King, L.A.L. & Elmhirst, R. (1914). Systematic investigation of a sandy shore. *Annual Report of the Scottish Marine Biological Association* 1914, 52–54.
- Kinne, O. (1971). Animals. Invertebrates. In: Kinne, O. (Editor). *Marine Ecology, Vol. I, Part 2*, pp. 821–995. Wiley Interscience, London.
- Little, C. & Kitching, J.A. (1996). *The Biology of Rocky Shores*. Oxford University Press, Oxford.
- McAllen, R., Taylor, A.C. & Davenport, J. (1998). Osmotic and body density response in the harpacticoid copepod *Tigriopus brevicornis* in supralittoral rockpools. *Journal of the Marine Biological Association of the United Kingdom* 78, 1143–1153.
<https://doi.org/10.1017/S0025315400044386>
- McAllister, S.M., Barnett, J.M., Heiss, J.W., Findlay, A.J., Macdonald, D.J., Dow, C.L., Luther III, G. W., Holly, A.M. & Chan, C.S. (2015). Dynamic hydrologic and biogeochemical processes drive microbially enhanced iron and sulfur cycling within the intertidal mixing zone of a beach aquifer. *Limnology and Oceanography* 60(1), 329–345.
<https://doi.org/10.1002/lno.10029>
- McIntyre, A.D. (1970). The range of biomass of intertidal sand, with special reference to the bivalve *Tellina tenuis*. *Journal of the Marine Biological Association of the United Kingdom* 50, 561–575.
<https://doi.org/10.1017/S0025315400004884>
- McLachlan, A. & Illenberger, W. (1986). Significance of groundwater nitrogen input to a beach/surf zone ecosystem. *Stygologia* 2, 291–296.
- Mauchline, J. (1967). The biology of *Schistomysis spiritus* (Crustacea, Mysidacea). *Journal of the Marine Biological Association of the United Kingdom* 47, 383–396.
<https://doi.org/10.1017/S0025315400056496>
- Millham, N.P. & Howes, B.L. (1994). Nutrient balance of a shallow coastal embayment. 1. Patterns of groundwater discharge. *Marine Ecology Progress Series* 112, 155–167.

- <https://doi.org/10.3354/meps112155>
- Montagna, P.A. & Yoon, W.B. (1991). The effect of freshwater inflow on meiofaunal consumption of sediment bacteria and microphytobenthos in San Antonio Bay, Texas, USA. *Estuarine, Coastal and Shelf Science* 33, 529–547.
[https://doi.org/10.1016/0272-7714\(91\)90039-E](https://doi.org/10.1016/0272-7714(91)90039-E)
- Moore, P.G. (2017). A prodigious collembolan 'swarming' event on Kames Bay, Millport. *The Glasgow Naturalist* 26(3), 86–87.
- Moore, P.G. & Beare, D.J. (1993). Taxonomic confusion in the genus *Pontocrates* (Crustacea: Amphipoda) and the presence of *P. arcticus* in Britain. *Journal of the Marine Biological Association of the United Kingdom* 73, 609–615.
<https://doi.org/10.1017/S0025315400033142>
- Morris, S. & Taylor, A.C. (1983). Diurnal and seasonal variation in physico-chemical conditions within intertidal rock pools. *Estuarine, Coastal and Shelf Science* 17, 339–355.
[https://doi.org/10.1016/0272-7714\(83\)90026-4](https://doi.org/10.1016/0272-7714(83)90026-4)
- Nicholls, A.G. (1935). Copepods from the interstitial fauna of a sandy beach. *Journal of the Marine Biological Association of the United Kingdom* 20, 379–405.
<https://doi.org/10.1017/S0025315400045306>
- Nicholls, A.G. (1939). Some new sand-dwelling copepods. *Journal of the Marine Biological Association of the United Kingdom* 23, 327–341.
<https://doi.org/10.1017/S0025315400013928>
- Paerl, H.W. (1997). Coastal eutrophication and harmful algal blooms: importance of atmospheric deposition and groundwater as new nitrogen and other nutrient sources. *Limnology and Oceanography* 42, 1154–1165.
https://doi.org/10.4319/lo.1997.42.5_part_2.1154
- Phillips, F.E. & Fleeger, J.W. (1985). Meiofauna meso-scale variability in two estuarine habitats. *Estuarine, Coastal and Shelf Science* 21, 745–756.
[https://doi.org/10.1016/0272-7714\(85\)90070-8](https://doi.org/10.1016/0272-7714(85)90070-8)
- Powell, R. & Moore, P.G. (1991). The breeding cycles of females of seven species of amphipod (Crustacea) in the Clyde Sea area. *Journal of Natural History* 25, 435–479.
<https://doi.org/10.1080/00222939100770291>
- Reid, D.M. (1930). Salinity interchange between sea-water in sand and overflowing fresh-water at low tide. *Journal of the Marine Biological Association of the United Kingdom* 26, 604–614.
<https://doi.org/10.1017/S0025315400072982>
- Robinson, C., Gibbes, B., Carey, H. & Li, L. (2007). Salt-freshwater dynamics in a subterranean estuary over a spring-neap tidal cycle. *Journal of Geophysical Research* 112(C9), 1–15.
<https://doi.org/10.1029/2006JC003888>
- Santos, P.J.P., Castel, J. & Souza-Santos, L.P. (1996). Seasonal variability of meiofaunal abundance in the oligo-mesohaline area of the Gironde estuary, France. *Estuarine, Coastal and Shelf Science* 43, 549–563.
<https://doi.org/10.1006/ecss.1996.0087>
- Smith, R.I. (1955). Salinity variation in interstitial water of sand at Kames Bay, Millport, with reference to the distribution of *Nereis diversicolor*. *Journal of the Marine Biological Association of the United Kingdom* 34, 33–46.
<https://doi.org/10.1017/S0025315400008596>
- Stephen, A.C. (1928). Notes on the biology of *Tellina tenuis* da Costa. *Journal of the Marine Biological Association of the United Kingdom* 15, 683–702.
<https://doi.org/10.1017/S0025315400009590>
- Stephen, A.C. (1929). Studies on the Scottish marine fauna. The fauna of the sandy and muddy areas of the tidal zone. *Transactions of the Royal Society of Edinburgh* 56, 291–306.
<https://doi.org/10.1017/S0080456800013211>
- Stephen, A.C. (1931). Notes on the biology of certain lamellibranchs on the Scottish coast. *Journal of the Marine Biological Association of the United Kingdom* 17, 277–300.
<https://doi.org/10.1017/S0025315400050840>
- Stephen, A.C. (1932). Notes on the biology of some lamellibranchs in the Clyde area. *Journal of the Marine Biological Association of the United Kingdom* 18, 51–68.
<https://doi.org/10.1017/S0025315400051298>
- Stephen, A.C. (1938). Production of large broods in certain marine lamellibranchs with a possible relation to weather conditions. *Journal of Animal Ecology* 7, 130–143.
<https://doi.org/10.2307/1119>
- Stephen, A.C. (1945). Molluscan ecology. *Annual Report of the Scottish Marine Biological Association* 1944–45, 14–16.
- Stephen, A.C. (1953). Life in some sandy shores. In: Marshall, S.M. & Orr, A.P. (Editors). *Essays in Marine Biology*, pp. 50–72. Oliver & Boyd, Edinburgh.
- Thomason, J.C., Hills, J. M. & Bennett, A. (1997). Dynamics of a massive freshwater influx in the Clyde Sea. *Estuarine, Coastal and Shelf Science* 44, 385–393.
<https://doi.org/10.1006/ecss.1996.0141>
- Warwick, R.M. (1989). The role of meiofauna in the marine ecosystem: evolutionary considerations. *Zoological Journal of the Linnean Society of London* 96, 229–241.
<https://doi.org/10.1111/j.1096-3642.1989.tb02258.x>
- Watkin, E.E. (1938). A revision of the amphipod genus *Bathyporeia* Lindström. *Journal of the Marine Biological Association of the United Kingdom* 23, 211–136.
<https://doi.org/10.1017/S0025315400054096>
- Watkin, E.E. (1939). The pelagic phase in the life history of the amphipod genus *Bathyporeia*. *Journal of the Marine Biological Association of the United Kingdom* 23, 467–481.
<https://doi.org/10.1017/S0025315400014028>
- Watkin, E.E. (1941). Observations on the night tidal migrant Crustacea of Kames Bay. *Journal of the Marine Biological Association of the United Kingdom* 25, 81–96.
<https://doi.org/10.1017/S0025315400014284>
- Watkin, E.E. (1942). The macrofauna of the intertidal sand of Kames Bay, Millport, Buteshire.

Transactions of the Royal Society of Edinburgh
60, 543–561.
<https://doi.org/10.1017/S0080456800008838>